<u>Math 115 - Test 2</u> October 16, 2014

Name _____ Score _____

Show all work to receive full credit. Supply explanations where necessary.

- 1. (10 points) Determine whether each statement is true (T) or false (F).
 - (a) _____ The coefficient of variation is used to measure the center of a data set.
 - (b) The 50th percentile is the same as the 3rd quartile.
 - (c) _____ All probabilities must be between 0 and 1.
 - (d) _____ About one-quarter of a data set falls below Q_1 .
 - (e) _____ A subjective probability is assigned by counting possible outcomes.
 - (f) _____ An outlier is any number above Q_3 or below Q_1 .
 - (g) _____ In a probability experiment, an event is any subset of the sample space.
 - (h) _____ The 2nd quartile of a data set is always equal to the mean.
 - (i) _____ It is impossible to have a z-score of zero.
 - (j) _____ If the probability of an event is close to 1, then the event is very likely.

2.	2. (5 points) State wheth	er each probability is theoretical, experimental, or subjective.
	, - ,	nt to drive in the snowstorm because she thought she had a 90%
	(b) The probability t	hat a flipped coin will land heads up is $1/2$.
	(c) In rolling a pair of rolling doubles	of dice 50 times, a double was rolled 10 times. The probability is $1/5$.
		ad driven through an intersection, there had been 37 accidents. f an accident is $37/1500$.
		in his pocket, and two of them are quarters. If he grabs a coin robability that it will be a quarter is $2/7$.
3. (3 points) Without attempting to compute anything, which do you think would greater: the coefficient of variation (CV) of adult American men's heights or the of their weights? Briefly explain why you think so?		


4. (10 points) Sammy Sosa was a major league baseball player from 1989 to 2007. His numbers of yearly regular season home runs are shown below in the order in which they occurred.

Compute the five-number summary, the interquartile range (IQR), and the cutoff values for outliers. Then sketch the boxplot on graph paper.

5. (3 points) For Yellowstone's Old Faithful geyser, the mean time between eruptions is 1.55 hr with a standard deviation of 0.11 hr. For Yellowstone's Lone Star geyser, the mean is 3.00 hr with a standard deviation of 0.16 hr. Compute the coefficient of variation (CV) for each geyser. Which geyser's eruption cycle has more variation?

6. (3 points) A test has mean score 73.4 and standard deviation 4.9. Sally scored 75 on the test, and Jake scored 72. Without actually computing their z-scores, state whose z-score is greater. Explain why you think so.

7. (3 points) The boxplot shown below describes a certain collection of data. Find approximate values for the five-number summary and the interquartile range (IQR).



8.	(6 points) Believe it or not, zenzizenizenic is an actual word. Suppose a letter is selected at random from this word.		
	(a) Give a possible sample space for this experiment.		
	(b) Are the outcomes in your sample space equally likely? Explain.		
	(c) Find the probability of each outcome in your sample space.		
	(d) Are the probabilities that you gave above above theoretical, experimental, or subjective?		

9. (4 points) The data below show the maximum daily air temperatures (in °F) measured at the DeKalb, IL weather station in January 2014.

19.2 1.5 18.8 30.3 20.7-6.92.2 8.525.437.2 36.438.7 38.9 34.419.7 34.0 15.6 19.3 41.1 35.0 10.5 14.54.427.933.0 30.1 1.5 3.5 21.530.817.1

(a) Arrange the temperatures in order from least to greatest.

(b) Find the percentile for 35°F.

(c) What temperature is at the 50th percentile?

10. (3 points) In January 2014, the mean maximum daily temperature was $21.8^{\circ}F$ with a standard deviation of $12.8^{\circ}F$. Compute the z-score for $-6.9^{\circ}F$. Do you think that $-6.9^{\circ}F$ was an unusually low temperature? Explain.